

德国 HYDRO-BIOS 公司--Ruttner 标准水样采集器

Standard Water Sampler acc. to Ruttner



这台便捷的多功能水样采集器(有机玻璃采水器)，可以从任何预期的深度采集水样。多年的实践已经证明，它所具有的闭锁器是很受欢迎的一款。这台水样采集器设计简单实用，并且确保可靠的采样结果。

这台标准水样采集器，系在绳索上，沉入水中，在这个过程中，采样器保持敞开。当到达预期的深度时，将使锤从绳索上释放，当它击中标准水样采集器时，闭锁器开始作用，采样管关闭。

一个量程为-2°C~+30°C的温度计，会显示样品的温度。温度可以很容易地透过水样采集器的透明塑料管读出。样品可以从水样采集器底盖的排水旋塞中排出，进行各种分析研究。

Ruttner 标准水样采集器(有机玻璃采水器)订购信息：

436 131 Ruttner 标准水样采集器

容积 1000ml，长 56cm，直径 10cm，质量 4.5kg

436 132 Ruttner 标准水样采集器

容积 2000ml，长 75cm，直径 10cm，质量 5kg

Ruttner 标准水样采集器(有机玻璃采水器)备用部件：

436 136 温度计

量程-2°C~+30°C，最小刻度 0.2°C，3 支/包

440 000 不锈钢使锤

孔径 6mm, 质量 400g

Ruttner 标准水样采集器照片集:

	
下水前的 Ruttner 标准水样采集器	采水结束的 Ruttner 标准水样采集器
	
采水结束的 Ruttner 标准水样采集器	处于关闭状态的 Ruttner 标准水样采集器

代表文献：

- 1.W. DAVISON,1977.Sampling and handling procedures for the polarographic measurement of oxygen in hypolimnetic waters.Freshwater Biology.7(4):393–401.
- 2.Austin B. M. Egbore,1978.Seasonal variations in the density of a small West African lake.Hydrobiologia.61(3):195-203.
- 3.Dr. U. Zaiss, P. Winter, H. Kaltwasser,1982.Microbial methane oxidation in the River Saar.Journal of Basic Microbiology.22(2):139–148.
- 4.V.F. Samanidou & I.N. Papadoyannis,1992.Study of heavy metal pollution in the waters of Axios and Aliakmon rivers in northern Greece.Journal of Environmental Science and Health . Part A: Environmental Science and Engineering and Toxicology.27(3):587-601.
- 5.Nilgün Kazancı, Reiner-Hartmut Plasa, Eike Neubert & Afife İzbırak,1992.On the limnology of Lake Köycegiz (SW Anatolia).Zoology in the Middle East.6(1):109-126.
- 6.Eduardo González-Mazo, Jesus María Forja, Abelardo Gómez-Parra ,1998.Fate and Distribution of Linear Alkylbenzene Sulfonates in the Littoral Environment.Environ. Sci. Technol..32(11):1636–1641.
- 7.V.M León, E González-Mazo, A Gómez-Parra,2000.Handling of marine and estuarine samples for the determination of linear alkylbenzene sulfonates and sulfophenylcarboxylic acids.Journal of Chromatography A.889(1-2):211–219.
- 8.Claus-Peter Stelzer,2001.RESOURCE LIMITATION AND REPRODUCTIVE EFFORT IN A PLANKTONIC ROTIFER.Ecology.82(9):2521–2533.
- 9.Udo Noack, Thomas Geffke, Ramani Balasubramanian, Jutta Papenbrock, Mike Braune, Dirk Scheerbaum,2004.Effects of the Herbicide Metazachlor on Phytoplankton and Periphyton Communities in Outdoor Mesocosms.Acta hydrochimica et hydrobiologica.31(6):482–490.
- 10.L. R. Rodríguez-Gallego, N. Mazzeo, J. Gorga, M. Meerhoff, J. Clemente, C. Kruk, F. Scasso, G. Lacerot, J. García, F. Quintans,2004.The effects of an artificial wetland dominated by free-floating plants on the restoration of a subtropical, hypertrophic lake.Lakes & Reservoirs: Research & Management.9(3-4):203–215.
- 11.Kristina Samuelsson, Johnny Berglund, and Agneta Andersson,2006.Factors structuring the heterotrophic flagellate and ciliate community along a brackish water primary production gradient.Journal of Plankton Research.28(4):345-359.
- 12.George Kehayias, Ekaterini Chalkia, Stavroula Chalkia, George Nistikakis, Ierotheos Zacharias, Anastasios Zotos,2008.Zooplankton dynamics in the upstream part of Stratos reservoir (Greece).Biologia.63(5):699-710.
- 13.MAJANEVA Markus, AUTIO Riitta, HUTTUNEN Maija, KUOSA Harri, KUPARINEN Jorma,2009.Phytoplankton monitoring: the effect of sampling methods used during different stratification and bloom conditions in the Baltic Sea.Boreal environment research.14(2):313-322.